OBSERVING REQUEST
University of Arizona Observatories

Year: 2015 **Term:** Jul–Dec **Proposal type:** short-term*

Photometric U and B Band Observations of Bright Transiting Exoplanets with the 61”/Mont4k

P.I.: M. Ryleigh Fitzpatrick (SO; *fitzpatrickm@email.arizona.edu*; 480-334-0880)

CoI(s): Caitlin Griffith (LPL), Zachary Watson (SO), Robert Zellem (JPL), Kyle Pearson (NAU), Jenny Calahan (SO), Quadry Chance (SO), Marina Dunn (SO), Andrew Henrici (SO), Nicolas Montiel (SO), Dominic Sanchez (SO), AzGOE Group (SO/LPL)

Abstract of Scientific Justification

Observations of transiting exoplanets yield their atmospheric temperature-pressure profiles, compositions, and even the presence of clouds/hazes and atmosphere circulation. However, interpreting measurements of transiting exoplanets is complicated by model degeneracies between the atmospheric pressure level and atmospheric composition; a $\sim$1% difference in the estimated planetary radius at the $\sim$1 bar pressure level can result in variations in molecular abundances of an order of magnitude or more. Here we request exoplanet atmosphere observations *outside* of the strong molecular and atomic absorption spectral regions, at optical wavelengths where the opacity is dominated by Rayleigh scattering and (if present) clouds/haze. We propose a concentrated observing campaign on 7 bright transiting exoplanets with the 61”/Mont4K in two optical filters (U and B). These targets have near-IR data, but lack measurements that are uncontaminated by strong molecular and/or atomic gas absorption features. Our 61” observations of XO-2b will be in support of a larger study to highly constrain this planet’s H_2O abundance with past and upcoming *Hubble*/WFC3 and Gemini data. We aim to disentangle the physical radius and the atmospheric composition of these exoplanets, as well as monitor these targets for the presence and potential variability of clouds. Our observations, coupled with radiative transfer modeling, will determine an upper limit to the radius of each planet at a specified pressure level by breaking the degeneracy between the assumed/derived radius and composition.

Summary of observing runs requested for this project

Run	Telescope	Cage	Instrument	PI	AO	Nights	Moon	Scheduling Optimal	Scheduling Acceptable	Sharing Poss.	Sharing Adv.
1	61”	f/13.5	Mont4k				no pref.	Jan–Jul	Jan–Jul	no	no

Scheduling constraints and unusable dates *(up to 4 lines)*: Please see “Time Requested” for details. We must observe during times that our exoplanet targets transit in front of their host stars. This restricts our requested time to specific nights and hours during those nights.

no text past this line

A $\star$ appended to the proposal type indicates a continuation proposal; a $\star$ appended to the name of a proposer indicates the proposer is a (graduate) student; a proposer whose name is underlined is certified on the proposed telescope/instrument combination; if a $*$ appears within the PI or AO box in the observations summary table, the instrument is a PI instrument and/or Adaptive Optics are requested – signatures are required on the next page.

Target list (attach list if longer than 26 objects)

#	Object	RA	Dec	mag / color / type / redshift / comment / etc.
1	HAT-P-1b	22:57:46.844	+38:40:30.33	V=9.87
2	HAT-P-32b	02:04:10.278	+46:41:16.21	V=11.44
3	HAT-P-33b	07:32:44.218	+33:50:06.12	V=11.03
4	HAT-P-41b	19:49:17.439	+04:40:20.724	V=11.36
5	WASP-12b	06:30:32.794	+29:40:20.29	V=11.57
6	WASP-33b	02:26:51.0582	+37:33:01.73	V=8.14
7	XO-2b	07:48:07.0	+50:13:33	V=11.18

Approval for Instrument Use from PI: N/A

Graduate students *(provide the following information for **each** student named as PI or CoI on the cover page. Have the advisor's signature(s) appear on **all** submitted copies)*

Student's Name	Advisor's Name	Advisor's Signature	2nd-yr	Thesis

Scientific Justification

Investigations of the atmospheric composition of transiting exoplanets have detected the main C and O-bearing species (H_2O, CO, CO_2 and CH_4) for a few of the brightest systems (e.g., Tinetti et al. 2007; Burrows et al. 2007; Knutson et al. 2007; Swain et al. 2008). During primary transit, when a planet passes in front of its host star, the light that transmits through its atmosphere reveals planetary absorption features. Transit observations include a fixed component of the solid disk of the planet and a wavelength-dependent component that is absorbed or scattered by the atmosphere. Such observations probe the $\sim$1 bar–1 millibar altitude region of the exoplanet's day-night terminator and scale with height. These measurements are directly sensitive to the column depth (at the exoplanet's limb) of individual molecular and atomic species and are only weakly dependent on temperature (for a given scale height).

The most limiting aspect in the analysis of exoplanet transmission observations is the degeneracy in the radius as a function of atmospheric pressure and composition (Benneke & Seager 2012, 2013; Griffith 2014). The derived pressure level at which the transit radii observations originate depends on the assumed abundance of molecular and atomic species in the atmosphere. Molecular abundances derived from primary transit spectra are constrained only to within $\sim$3–5 orders of magnitude (e.g., Madhusudhan & Seager 2009). The derived abundances' dependence on the radius is significant: a $\sim$1% difference in the estimated planetary radius of a 1000 K hot Jupiter at the $\sim$10 bar pressure level can result in variations in molecular abundances of 2 orders of magnitude or more (Griffith 2014). Essentially the solution set is degenerate; model atmospheres can have either a large radius and low gas mixing ratio or smaller radius and high gas mixing ratio, and block the host stars' light equally well.

This degeneracy can be further complicated by the existence of clouds. Recent studies at visible wavelengths have found that some exoplanets have clouds (e.g., Demory et al. 2013, Kreidberg et al. 2014, Knutson et al. 2014), while others have relatively clear atmospheres (e.g., Fraine et al. 2014). However, it is unclear why some planets have clouds while others do not. Clouds can potentially indicate the planet's rotation rate for relatively fixed and broad-coverage clouds (Seager 2010) and influence the interpretation of both photometric and spectroscopic data.

We propose a concentrated, investigative study of the optical spectral characteristics of 8 exoplanets at wavelengths (0.35–0.55μm) that lie outside the strongest molecular gas absorption signatures. Here, the atmospheric opacity is dominated by H_2 Rayleigh scattering and, if present, clouds. Thus, we can disentangle the degeneracy between the physical radius and the atmospheric composition of the exoplanets by probing these "clear" spectral regions in which we can confidently assume absorption due to H_2 alone. In this way, we are able to constrain the radius of the hot Jupiter XO-2b to 56σ using 61"/Mont4k U and B-band photometric data (Fig. 1; Griffith 2014; Zellem et al. 2015). In addition, we will be able to conduct a dedicated study to detect (via the slope of our visible data, which will deviate from a H_2 Rayleigh scattering slope) and monitor possible cloud coverage in transiting exoplanet atmospheres.

This proposal concentrates on 7 of the brightest (V >11.25) exoplanets, offering the ability to conduct high precision measurements. Some of them have large radii, and therefore high contrast, while others have a large scale height, and therefore a large transit signal. They already have a substantial amount of data longward of$\sim$0.60 μm (at least 5 photometric points), to which we can anchor our bluer data. We will measure the primary transit depths of these exoplanets in the B ($\sim$0.34–0.54 μm) and U ($\sim$0.305–0.425 μm) bands, which are free of any molecular or atomic absorption features.

For our 61"/Mont4k targets (Table 1), our observations will be the bluest observations to date of their transmission spectra, and the only observations outside of strong molecular or atomic absorption bands. The data will be reduced and analyzed to determine first whether the slope of optical absorption in the atmospheres of exoplanets generally mimics that of Rayleigh scattering (see Fig. 2). Such a finding, coupled with a lack of temporal variability, would suggest atmospheres devoid of clouds. This outcome would also allow us to derive an upper limit on the radius of the planet, as the opacity would be due to Rayleigh scattering of H_2 only, and the existence of clouds would imply a smaller radius. We will model the optical and published near-IR observations to constrain H_2O and potentially CO in the exoplanetary atmospheres given our constraints for the radius.

In particular, our 61" observations of XO-2b are in support of a larger program that aims to determine the C and O abundances to within a factor of 3 with coordinated, past, and upcoming Gemini/GMOS and Hubble/WFC3 observations (PI: C. A. Griffith). With the 61", we will measure where XO-2b's atmospheric opacity is dominated by Rayleigh scattering and, if present, clouds (Fig. 2). Thus we can disentangle the degeneracy between the physical radius and the atmospheric composition of the exoplanet by probing these clear spectral regions in which we can confidently assume absorption due to H_2 alone. However high-precision observations at blue wavelengths are difficult because of high telluric extinction and telescope systematics. Here we propose to treat the resulting errors with simultaneous observations of XO-2b with Mayall/KOSMOS, the Discovery Channel Telescope (DCT) in northern Arizona, and the 61" telescope. Since the transit depth will be a common signal shared between these telescopes, we will run a joint analysis of all datasets (see Experimental Design) to remove the non-common signals, i.e., red noise sources, and get closer to the photon noise limit. This analysis will also allow us to characterize the systematics of each platform so they can be removed in future studies.

This project also includes participation from UofA astronomy and physics undergraduate members of the Arizona Ground-based Observing of Exoplanets group (AzGOE). This group uses UofA telescopes to characterize the thermal structure and molecular composition of transiting exoplanets while stressing undergraduate involvement in all parts of the analysis. PI M. Ryleigh Fitzpatrick and Co-I Zachary Watson, under the guidance of Co-I Caitlin Griffith, will lead this group of students as they learn how to observe transiting exoplanets and then reduce the data. The younger students will be involved in preparing and conducting observations and performing data reduction and analysis, whereas the older students will be involved in preparing observation proposals, learning the syntax for formal scientific publications, and the application of our results to the study of exoplanet atmospheres. Since extrasolar planets are at the forefront of observational astronomy, AzGOE provides the opportunity for participating members to be on high impact papers, as well as gain valuable experience in this field of research.

References

Benneke, B., & Seager, S. 2012, ApJ, 753
Benneke, B., & Seager, S. 2013, ApJ, 778
Burrows, A., et al. 2007, ApJL, 668, 171L
Crouzet, N., et al. 2013, arXiv
Demory, B.-O., et al. 2013, arXiv, 1309.7894
Dittmann, J., et al. 2009, ApJ, 701, 756
Dittmann, J., et al. 2010, ApJ, 717, 235
Fraine, J., et al. 2014, Nature , 513, 526
Griffith, C. A. 2014, Royal Soc., 372
Kipping, D., et al. 2014, ApJ , 784, 28
Knutson, H., et al. 2007, Nature, 447, 183
Knutson, H., et al. 2014, Nature , 505, 66
Kreidberg, L. et al. 2014, Nature , 505, 69
Madhusudhan, N., & Seager, S. 2009, ApJ, 707, 24
Madhusudhan, N., et al. 2011, Nature, 469, 64
Seager, S. 2010, Exopl. Atmos., Princeton U. Press
Swain M. R., et al. 2008, Nature, 452, 329
Swain, M. R. et al. 2009b, ApJ, 704, 1616
Swain, M. R., et al. 2009a, ApJL, 690, L114
Swain, M. R., et al. 2010, Nature , 463, L
Teske, J. et al. 2013, MNRAS, 431, 1669
Thatte, A. et al. 2011, A&A, 523, A35
Tinetti, G., et al. 2007, Nature, 448, 169
Tinetti, G., et al. 2010, ApJ , 712, L139
Turner, J. D, et al. 2013, MNRAS, 428, 678
Waldmann, I. P., et al. 2012, ApJ , 744, 35
Waldmann, I. P., et al. 2013, ApJ , 766, 7
Zellem, R. T. et al. 2014, ApJ , 796, 48
Zellem, R. T. et al. 2015, in prep



Figure 1: A sample B-filter lightcurve of XO-2b which demonstrates the quality we are capable of achieving with the 61"/Mont4k. Its associated best-fit model is over plotted in red and the corresponding residuals are in the bottom frame. From such a lightcurve, we can make a 53σ detection of XO-2b's primary transit ($R^2_{planet}/R^2_{star} = 0.0106 \pm 0.0002$). (Zellem et al. 2015)



Figure 2: A plot of our 61"/Mont4k U and B-band photometry (points shortward of 0.6 μm) and previous $Hubble$/NICMOS measurements for XO-2b (all other data; Crouzet et al. 2013) compared to radiative transfer atmospheric models that include Rayleigh scattering, pressure-induced H_2 absorption with different 10 bar optically-thick planetary radii (R_{10}) and water abundances ([H_2O]). Our current retrieval of XO-2b's radius is largely limited by the scatter in our B-band photometry. We propose to observe XO-2b with additional nights on the 61"/Mont4k here not only to better constrain its radius via reproducibility but also to determine if its varying B-band radius is due to stellar variability or platform-specific systematics. (Griffith 2014)

Experimental Design & Technical Description *Describe your overall observational program. How will these observations contribute toward the accomplishment of the goals outlined in the science justification? If you've requested long-term status, justify why this is necessary for successful completion of the science.* ***(up to one page)***

We request 14 nights to observe the primary transits of 7 different exoplanets (see Table 1) using the 61"/Mont4K. The Mont4k CCD on the 61" telescope has been shown to produce accurate photometry in transit-observation campaigns on other targets (Dittmann et al. 2009; Dittmann et al. 2010; Turner et al. 2013; Teske et al. 2013; Griffith 2014; Zellem et al. 2015). This campaign is a continuation of TBS-proposals from 2012B and 2013A, UAO-S64 from 2013A, UAO-S58 from 2013B,UAO-S60 from 2014A, UAO-S86 from 2014B, and UAO-S225 from 2015A.

Obtaining photometry or spectroscopy across $\sim$0.3–0.5 μm is necessary to constrain the slope of the optical transmission spectrum of these bright exoplanets and fulfill our scientific objective. To determine to 1σ-confidence if the short-wavelength ($<$0.50 μm) slope is Rayleigh scattering (Fig. 2), we need an error on our transit-depth measurement of $\sim$1% or less. Previously with 61"/Mont4k (Fig. 1), we obtained errors of $\sim$1% in U-band with one night of good observing (Griffith 2014; Zellem et al. 2015). However, while we only need one good observation to obtain the required precision for this project, we are requesting at least two per target to confirm our observations and in case of host star variability or weather. Any residual systematics will be treated with an array of procedures that we have successfully used to extract the planetary signal, such as the Self-Coherence Method, Independent Component Analysis (ICA), and Principal Component Analysis (PCA) (e.g., Swain et al. 2010, Thatte et al. 2010, Waldmann et al. 2012, Waldmann et al. 2013, Zellem et al. 2014). Our 61" data on XO-2b will also be compared with that measured by the DCT and Mayall and analyzed with an ICA and PCA to determine whether features detected in the light-curve are red noise specific to the observing platform intrinsic to the planetary system. In this way, we will quantify the effect of systematics specific to each observing platform, which will help future studies remove their effects and achieve a higher SNR.

We need to observe full transits, as well as pre- and post-transit baseline, using both filters. To optimize SNR levels and get faster readout times, we can use 3x3 binning of the CCD if necessary for the fainter target in our sample. We will be using the Exoplanet Data Reduction Pipeline to reduce our data (Turner et al. 2013; Teske et al. 2013; Zellem et al. 2015) and an already-developed in-house modeling package to obtain the planetary parameters.

A large portion of the 61" photometry data reduction will be conducted by Steward Observatory undergraduate members of the Arizona Ground-based Observing of Exoplanets group (AzGOE), under direction of PI M. Ryleigh Fitzpatrick and Co-I Zachary Watson, who will work closely with Co-I Caitlin Griffith. The radiative transfer analysis and the implications for the planet's physical characteristics will be conducted by Co-I Caitlin Griffith.

These are time-sensitive observations, as there is very limited data in the transmission spectrum of our targets $\lesssim$0.50 μm. Our observations will be the bluest of these targets to date. We will publish this data detailing our results and place them in context with the existing optical and NIR observations. We will constrain the exoplanets' radii, which is essential to enabling accurate and informative modeling of their atmospheric compositions. Previous 61" photometric data of XO-2b (Fig. 1) has allowed us to successfully constrain its radius and, when coupled with radiative transfer modeling of this data and other published observations, its molecular composition (Griffith 2014; Zellem et al., 2015).

Summary of Time Requested and Awarded *The TAC needs to understand the scope of this project — (1) tell us how many UAO nights you've already had for this project, how many you request this time, and (a good guess of) how many you need to complete the project; (2) if a substantial amount of observing for this project comes from non-UAO telescopes, tell us about that observing, and how the UAO part fits in; (3) if you are collaborating with people who have telescopes, especially if you are part of a large collaboration, tell us who is leading the project, and how UAO time and your participation fit in.* ***(up to one page)***

We were awarded 4 TBS nights on the Kuiper 61" UAO telescope in 2012B and 3 nights in 2013A for this project on XO-2b. The results from these observations are to be published shortly. We were awarded 10 TBS nights during the summer of 2013A; however, due to monsoons, all but 1 of the nights were lost. We were rewarded with 8 nights for semester 2013B, 3 of which were lost due to weather. We were awarded 11 nights for the semester 2014A; all but 1 were lost due to weather. We were awarded 6 nights for last semester (2014B) and 13 nights this semester (2015A).

The following table lists observing opportunities for our selected bright exoplanet targets at the 61" for 2015B.

Note: We are requesting 2 nights per target. In the table below, we list each night in order of preference, with the top 2 nights notated in bold. However, our observations of XO-2b are part of a larger collaboration, including *Hubble*/WFC3, Gemini, Mayall 4 m, and Discovery Channel Telescope time. We are requesting 4 nights here in support of this large-scale study.

Table 1: **Requested Primary Transit Dates & Times**

Target	**Date Requested** local start night	**Ingress** local time	**Egress** local time	**Platform**
HAT-P-1b	13-Sept	22:24	01:11	61"/Mont4k
HAT-P-1b	04-Sept	00:04	02:51	61"/Mont4k
HAT-P-1b	26-Aug	01:44	04:31	61"/Mont4k
HAT-P-32b	24-Oct	22:49	01:55	61"/Mont4k
HAT-P-32b	21-Nov	21:37	00:43	61"/Mont4k
HAT-P-32b	26-Sept	00:00	03:07	61"/Mont4k
HAT-P-32b	08-Nov	00:01	03:07	61"/Mont4k
HAT-P-33b	01-Dec	00:38	05:03	61"/Mont4k
HAT-P-41b	03-Aug	21:39	01:44	61"/Mont4k
WASP-12b	03-Dec	00:16	03:11	61"/Mont4k
WASP-12b	15-Dec	00:24	03:19	61"/Mont4k
WASP-12b	26-Dec	22:20	01:16	61"/Mont4k
WASP-12b	27-Dec	00:32	03:27	61"/Mont4k
WASP-33b	01-Nov	22:30	01:12	61"/Mont4k
WASP-33b	23-Nov	21:29	00:11	61"/Mont4k
WASP-33b	10-Oct	23:31	02:13	61"/Mont4k
WASP-33b	21-Oct	23:01	01:42	61"/Mont4k
XO-2b	23-Dec	01:51	04:32	61"/Mont4k
XO-2b	10-Dec	23:57	02:38	61"/Mont4k
XO-2b	10-Nov	01:42	04:23	61"/Mont4k
XO-2b	31-Dec	22:11	00:52	61"/Mont4k

Previous Use of Steward Facilities *List **all** allocations of telescope time for the present project and allocations for other projects on facilities available through UAO during the past 2 years, together with the current status of the data (cite publications where appropriate). Mark those allocations related to the present proposal (i.e, precede text with* `\related` *command). (**up to one page**)*

- ⋆ Awarded 4 TBS nights in 2012B and 3 nights in 2013A (UAO-S64) for observations of XO-2b using the Kuiper 61". These observations have been reduced.
- ⋆ Awarded 10 TBS nights in 2013B for observations for the exoplanet study on the 61"/Mont4k. All but 1 night were lost to monsoons. These observations have been reduced.
- ⋆ Awarded 8 nights on the 61"/Mont4k to continue this exoplanet study in 2013B (UAO-S58). 3 nights were lost due to weather. These observations have reduced.
- ⋆ Awarded 11 nights on the 61"/Mont4k this semester (2014A), all but 1 were lost due to weather. Also awarded 2 nights on VATT/VATTSpec (1 for training + 1 for observing). We are currently reducing and analyzing data. Some of this data is included in Zellem et at. 2015, in prep.
- ⋆ Awarded 4 nights last semester (2014B) on the 61"/Mont4k and 2 nights on VATT/VATTSpec. We are currently reducing and analyzing data. Some of this data is included in Zellem et at. 2015, in prep.
- ⋆ Awarded 13 nights this semester (2015A) on the 61"/Mont4k and 2 nights on the VATT/VATTSpec. Currently observing and reducing. Some of this data is included in Zellem et at. 2015, in prep.